

Osteoporosis Management in Women with a Fracture– Cheat Sheet

Weight: 1	Current Star: 2	Current Standing: 39%	Next Threshold: 41%
------------------	------------------------	------------------------------	----------------------------

Star Measure Description
This measure shows the percent of female plan members who broke a bone and got screening or treatment for osteoporosis within 6 months.

Measure Source
<ul style="list-style-type: none"> • Chart Review & Claims • 7/1/-6/30

Requirement Specifications
The percentage of woman MA enrollees 67 - 85 who suffered a fracture and who had either a bone mineral density (BMD) test or prescription for a drug to treat osteoporosis in the six months after the fracture.

Best Practice/ Call to Action
<ul style="list-style-type: none"> • Recommend/schedule DXA scan for those with recent qualifying fracture or prescribe qualifying osteoporosis treatment medications within 6 months of the fracture • Medical Record alerts (for clinics that have capability) for patients that had a fracture

Medications
<ul style="list-style-type: none"> • Biphosphonates • Calcitonin • Denosumab • Raloxifene • Teriparatide

Star Ratings Performance & Trends					
Year	1-Star	2-Star	3-Star	4-Star	5-Star
2020	< 31%	≥ 31% to < 41%	≥ 41% to < 50%	≥ 50% to < 67%	≥ 67%
2019	<29%	≥29% to <45%	≥45% to <57%	≥57% to <78%	≥ 78%
2018	<24%	≥24% to <42%	≥42% to <52%	≥52% to <71%	≥ 71%
2017	<21%	≥21% to <34%	≥34% to <51%	≥51% to <70%	≥70%