

Statin Therapy for Patients with Cardiovascular Disease – Cheat Sheet

Weight: 1	Current Star: 4	Current Standing: 83%	Next Threshold: 87%
------------------	------------------------	------------------------------	----------------------------

Star Measure Description

This measure shows the percent of plan members with heart disease who get the right type of cholesterol-lowering drugs.

Measure Source

- Prescription Drug Event data from CMS
- 1/1 – 12/31

Requirement Specifications

The percentage of males 21–75 years of age and females 40–75 years of age during the measurement year, who were identified as having clinical atherosclerotic cardiovascular disease (ASCVD) and were dispensed at least one high or moderate-intensity statin medication during the measurement year.

Best Practice/ Call to Action

- Patients with the following should be on a statin:
 - Any form of clinical ASCVD
 - Primary LDL-C levels of 190mg per dL or greater
 - Diabetes mellitus, 40 to 75 years of age, with LCL level of 70 to 189mg per dL
- Without diabetes, 40 to 75 years of age, with an estimated 10- year ASCVD risk of at least 7.5%

Exclusions

Must be coded annually:

- **G72.0:** Drug Induced myopathy
- **G72.2:** Myopathy due to other toxic agents
- **G72.9:** Myopathy, unspecified
- **M62.82:** Rhabdomyolysis
- **M79.1:** Myalgia

Medications

- Moderate or high dose statin

Star Ratings Performance & Trends

<u>Year</u>	<u>1-Star</u>	<u>2-Star</u>	<u>3-Star</u>	<u>4-Star</u>	<u>5-Star</u>
2020	< 75%	≥ 75% to < 79%	≥ 79% to < 83%	≥ 83% to < 87%	≥ 87%
2019	< 70%	≥70% to <76%	≥76% to <81%	≥81% to <85%	≥ 85%