



Statin Use in Persons with Diabetes (SUPD) – Cheat Sheet

Weight: 1	Current Star: 3	Current Standing: 80%	Next Threshold: 81%
------------------	------------------------	------------------------------	----------------------------

Star Measure Description
This measure shows the percent of plan members with diabetes who take the most effective cholesterol-lowering drugs.

Measure Source
<ul style="list-style-type: none"> NDC Lists (Prescription Drug Event data from CMS) 1/1 – 12/31

Requirement Specifications
This measure is defined as the percent of Medicare Part D beneficiaries 40-75 years old who were dispensed at least two diabetes medication fills who received a statin medication fill during the measurement period.

Best Practice/ Call to Action
<ul style="list-style-type: none"> Prescribe a statin for patients diagnosed with diabetes.

Exclusions
<p>Must be coded annually:</p> <ul style="list-style-type: none"> G72.0: Drug-induced myopathy G72.2: Myopathy due to other toxic agent G72.9: Myopathy, unspecified K74.60: Unspecified cirrhosis M60.9: Myositis

Star Ratings Performance & Trends						
<u>Year</u>	<u>Cut Points</u>	<u>1-Star</u>	<u>2-Star</u>	<u>3-Star</u>	<u>4-Star</u>	<u>5-Star</u>
2020	MA-PD	< 74%	≥ 74% to < 78%	≥ 78% to < 81%	≥ 81% to < 83%	≥ 83%
2019	MA-PD	< 72%	≥72% to <76%	≥76% to <80%	≥80% to <83%	≥ 83%