

### **Statin Use in Persons with Diabetes (SUPD) – Cheat Sheet**

<b>Weight: 1</b>	<b>Current Star: 3</b>	<b>Current Standing: 80%</b>	<b>Next Threshold: 81%</b>
------------------	------------------------	------------------------------	----------------------------

<b>Star Measure Description</b>
This measure shows the percent of plan members with diabetes who take the most effective cholesterol-lowering drugs.

<b>Measure Source</b>
<ul style="list-style-type: none"> <li>NDC Lists (Prescription Drug Event data from CMS)</li> <li>1/1 – 12/31</li> </ul>

<b>Requirement Specifications</b>
This measure is defined as the percent of Medicare Part D beneficiaries 40-75 years old who were dispensed at least two diabetes medication fills who received a statin medication fill during the measurement period.

<b>Best Practice/ Call to Action</b>
<ul style="list-style-type: none"> <li>Prescribe a statin for patients diagnosed with diabetes.</li> </ul>

<b>Star Ratings Performance &amp; Trends</b>						
<u>Year</u>	<u>Cut Points</u>	<u>1-Star</u>	<u>2-Star</u>	<u>3-Star</u>	<u>4-Star</u>	<u>5-Star</u>
2020	MA-PD	< 74%	≥ 74% to < 78%	≥ 78% to < 81%	≥ 81% to < 83%	≥ 83%
2019	MA-PD	< 72%	≥ 72% to < 76%	≥ 76% to < 80%	≥ 80% to < 83%	≥ 83%