

Eye Exam – Cheat Sheet

Weight: 1	Current Star: 3	Current Standing: 70%	Next Threshold: 78%
------------------	------------------------	------------------------------	----------------------------

Star Measure Description

This measure shows the percent of members with diabetes who had retinal eye exam to check for diabetic retinopathy.

Measure Source

- Chart Review & Claims
- 1/1 – 12/31

Requirement Specifications

The percentage MA members between 18 and 75, with diabetes, (type 1 and type 2) who had an eye exam (retinal) performed during the measurement year.

Best Practice/ Call to Action

- Patients with type 2 diabetes should have an initial dilated and comprehensive eye examination by an ophthalmologist or optometrist at the time of the diabetes diagnosis.
- If there is no evidence of retinopathy for one or more annual eye exams, then exams every two years may be considered.
- Refer patients to ophthalmology or optometry for dilated retinal exam
- May refer for Retinal Camera.

Supplemental Data

CPT II Codes:

- **2022F:** Dilated eye exam w/interpretation by an ophthalmologist or optometrist documented and reviewed
- **2024F:** Seven standard field stereoscopic photos with interpretation by an ophthalmologist or optometrist documented and reviewed
- **2026F:** Eye imaging validated to match diagnosis from seven standard field stereoscopic photos results documented and reviewed
- **3072F:** Low risk for retinopathy

Star Ratings Performance Trends & Cut Points

Year	1-Star	2-Star	3-Star	4-Star	5-Star
2020	< 63%	≥ 63% to < 69%	≥ 69% to < 73%	≥ 73% to < 78%	≥ 78%
2019	< 56%	≥56% to <64%	≥64% to <73%	≥73% to <80%	≥ 80%
2018	< 47%	≥47% to <59%	≥59% to <72%	≥72% to <81%	≥ 81%



Administered by Health Alliance

3310 Fields South Drive
Champaign, IL 61822
OSFMedAdvantage.org

2017	< 46%	≥46% to <61%	≥61% to <73%	≥73% to <81%	≥ 81%
------	-------	--------------	--------------	--------------	-------